

# daflon<sup>®</sup> 500 mg

Micronized purified flavonoid fraction

## Presentation and composition

Box of 30 film-coated tablets.

Micronized purified flavonoid fraction corresponding to:

- diosmin: 90 %
- flavonoids expressed as hesperidin: 10 %

Excipients:

Core: gelatine, magnesium stearate, microcrystalline cellulose, sodium starch glycolate, talc.

Film-coating: glycerol, macrogol 6000, magnesium stearate, methylhydroxypropylcellulose, red iron oxide, sodium lauryl sulfate, titanium dioxide, yellow iron oxide.

## Marketing Authorisation Holder



Les Laboratoires Servier  
22, rue Garnier  
92200 Neuilly-sur-Seine  
France

## Manufacturer

Les Laboratoires Servier Industrie  
905, route de Saran  
45520 Gidy - France

## Therapeutic indications

This medicine is a venotonic (it increases venous tone) and a vasculoprotector (it increases resistance in small blood vessels).

It is recommended for treating venous circulation disorders (swollen legs, pain, restless legs) and for treating symptoms due to acute hemorrhoidal attack.

## Side effects

Some cases of routine gastric disorders and neurovegetative disorders (feeling of discomfort) have been reported, not requiring discontinuation of treatment.

*Report any untoward or undesirable effect which has not been mentioned in this package insert to your doctor or your pharmacist.*

### **Precautions**

#### *Acute hemorrhoidal attack:*

If the hemorrhoid symptoms do not disappear within 15 days, you should ask your doctor for advice.

#### *Venous circulation disorders:*

The most effective way of giving this treatment is in combination with a healthy lifestyle. Avoid exposure to the sun, heat, excessive standing and being overweight. Walking and wearing special support stockings stimulate blood circulation.

*If you are in any doubt, do not hesitate to ask your doctor or your pharmacist for advice.*

### **Interactions with other medicines and other forms of interactions**

*You must always tell your doctor or your pharmacist about any other treatment you are receiving.*

### **Pregnancy - Breast-Feeding**

Inform your doctor if you become pregnant.

In the absence of data on entering breast milk, breast-feeding should be avoided during treatment.

*In general, if you are pregnant or breast-feeding, you should always ask your doctor or pharmacist for advice before using a medication.*

### **Dosage and method of administration**

Oral route.

The tablets should be taken at meal times.

#### *Venous insufficiency:*

2 tablets daily, one at midday and one in the evening.

#### *Acute hemorrhoidal attack:*

A 4-day course of 6 tablets daily, followed by 4 tablets daily over the next 3 days.

*In all cases strictly comply with your doctor's prescription.*

### **Storage conditions**

Store below 30°C.

Do not use after the expiry date indicated on the outer packaging.

### **Marketing authorisation number**

MA 066/00501

### **Revision date of the leaflet**

March 2006